ARTICLE

<https://doi.org/10.1038/s41467-019-13251-w>

OPEN

Symmetry of molecular Rydberg states revealed by XUV transient absorption spectroscopy

Peng Peng^{1*}, Claude Marceau¹, Marius Hervé¹, P.B. Corkum¹, A.Yu. Naumov¹ & D.M. Villeneuve^{1*}

Transient absorption spectroscopy is utilized extensively for measurements of bound- and quasibound-state dynamics of atoms and molecules. The extension of this technique into the extreme ultraviolet (XUV) region with attosecond pulses has the potential to attain unprecedented time resolution. Here we apply this technique to aligned-in-space molecules. The XUV pulses are much shorter than the time during which the molecules remain aligned, typically <100 fs. However, transient absorption is not an instantaneous probe, because long-lived coherences re-emit for picoseconds to nanoseconds. Due to dephasing of the rotational wavepacket, it is not clear if these coherences will be evident in the absorption spectrum, and whether the properties of the initial excitations will be preserved. We studied Rydberg states of N₂ and O₂ from 12 to 23 eV. We were able to determine the polarization direction of the electronic transitions, and hence identify the symmetry of the final states.

¹Joint Attosecond Science Laboratory, National Research Council and University of Ottawa, Ottawa, ON K1A 0R6, Canada. *email: pengpengnrc@gmail.com; david.villeneuve@uottawa.ca

XUV transient absorption spectroscopy uses coherent extreme ultraviolet (XUV) pulses to probe the electronic structure of atoms and molecules. Attosecond transient absorption spectroscopy (ATAS) uses high harmonic generation to create a broad spectrum in the XUV or soft X-ray range. The broad spectrum permits the recording of many absorption lines in parallel^{1–17}. ATAS can probe valence electrons through excitation to high-lying electronic states^{1–14}. ATAS can also excite core-level electrons to reveal molecular structure using techniques such as near-edge X-ray absorption fine structure (NEXAFS)^{15–18}, for example around the carbon K-edge at 284 eV. Pump-probe type experiments in which two-photon processes are driven by a combination of an attosecond pulse and an infrared pulse have revealed coupling between states and the production of light-induced states^{4,7}. Shifts in absorption features have been used to observe vibrational motion in bromine molecules¹⁴.

When linear molecules are irradiated by an infrared pump pulse whose duration is much shorter than the rotational period, nonadiabatic field-free alignment is achieved^{19–21}. Each initial thermally-populated rotational eigenstate $|J_0, M_0\rangle$ will expand to a rotational wave packet.

$$\Psi_{J_0, M_0}(t) = \sum_{J, M} a_{J, M}^{J_0, M_0}(t) |J, M\rangle, \quad (1)$$

where J is the rotational quantum number, M is the projection on the polarization axis and $a_{J, M}^{J_0, M_0}(t)$ is the amplitude for each $|J, M\rangle$ state. The time evolution of the wave packet can be calculated by solving the time-dependent Schrödinger equation. The degree of alignment is characterized by:

$$\langle \cos^2\theta \rangle_{J_0, M_0}(t) = \langle \Psi_{J_0, M_0}(t) | \cos^2\theta | \Psi_{J_0, M_0}(t) \rangle, \quad (2)$$

in which θ is the angle of the molecular axis to the polarization axis. Finally, thermally averaged values of the degree of alignment are obtained as:

$$\langle \cos^2\theta \rangle(t) = \sum_{J_0, M_0} \rho_{J_0, M_0}(T) \langle \cos^2\theta \rangle_{J_0, M_0}(t), \quad (3)$$

where T is the rotational temperature and $\rho_{J_0, M_0}(T)$ is the Boltzmann weight function for the $|J_0, M_0\rangle$ initial states. $\langle \cos^2\theta \rangle(t)$ varies with time after the aligning laser pulse as various rotational states go in and out of phase. The probe attosecond pulses arrive at a revival of the rotational wavepacket at which the value of $\langle \cos^2\theta \rangle$ maximizes. The electronic excitation is in the molecular frame, but observations occur in the laboratory frame; the frames are related through the Wigner rotation matrix. The free induction decay resulting from the induced polarization in the molecule frame continues to radiate while the rotational wavepacket dephases. At times the radiation associated with different rotational eigenstates will be out of phase and will lead to cancellation of the emission. In addition, the rotational constants of the lower and upper states will be different. Therefore it needs to be demonstrated that the time-integrated absorption of aligned molecules will exhibit the characteristics of the attosecond excitation.

In the following, we use XUV transient absorption spectroscopy to record the molecular absorption spectra of N_2 and O_2 . The short pulse duration of attosecond pulse trains (typically 5–30 fs, but as short as 50 as refs. 22,23) permits using a near-infrared (NIR) laser pulse to impulsively align the molecular sample^{19–21}, allowing the absorption measurements to be made in a sample in which the molecular axes can be aligned relative to the polarization axis of the probing XUV field. This is not possible in synchrotron experiments due to their relatively long pulse durations. We will show that we can discriminate between parallel and perpendicular transitions for linear molecules, and hence

identify the symmetry of the final states. In the case of O_2 , the observed symmetry of some states differs from previous assignments. This same approach can be applied to larger molecules, for example asymmetric top molecules²⁴.

Results

XUV transient absorption in N_2 and O_2 . Figure 1a illustrates the experimental arrangement. A 10 fs NIR pulse centered at 800 nm was focused into a pulsed gas jet of xenon for high harmonic generation (see Methods for more details). The bandwidth of each harmonic was 2.3 eV and the total spectrum spanned 12–24 eV, as shown by the black curve in Fig. 1b. A 70 fs NIR pulse was used to perform impulsive alignment on the sample molecules. It was combined with the XUV pulse with a holey mirror. The alignment pulse was displaced from the center of the holey mirror while the XUV pulse passed through the hole. Both the XUV and the alignment pulses were vertically polarized, they propagated noncollinearly and intersected at the absorption gas jet with a crossing angle of 18 mrad. The transmitted XUV spectra were recorded at different time delays τ between the alignment and XUV pulses. The blue (red) curve in Fig. 1(b) shows the transmitted XUV spectrum at aligned (anti-aligned) delay $\tau = 8.57$ ps (anti-aligned delay $\tau = 8.32$ ps) for N_2 . About 10,000 laser shots were accumulated at each delay point.

We define the differential absorption spectrum to be the differential optical density $\Delta OD(E, \tau) = -\log_{10}[I_{\text{on}}(E, \tau)/I_{\text{off}}(E, \tau)]$. Here, $I_{\text{on}}(E, \tau)$ and $I_{\text{off}}(E, \tau)$ represent the XUV spectra after transmission through the sample, taken with and without the alignment field, respectively. This is effectively the XUV absorption of the aligned medium minus that of the randomly aligned medium. Fig. 2a shows the static absorption of isotropic N_2 measured by blocking the NIR alignment pulse. The ground state configuration may be written as $(1\sigma_g)^2(1\sigma_u)^2(2\sigma_g)^2(2\sigma_u)^2(1\pi_u)^4(3\sigma_g)^2 X^1\Sigma_g^+$. The adiabatic ionization energies corresponding to different final states of N_2^+ are $(3\sigma_g)^{-1}X^2\Sigma_g^+ = 15.58$ eV, $(1\pi_u)^{-1}A^2\Pi_u = 16.70$ eV, $(2\sigma_u)^{-1}B^2\Sigma_u^+ = 18.75$ eV^{25–27}, as shown by the thick gray lines in Fig. 2a. The XUV pulse excites numerous overlapping vibrational series associated with states of Rydberg and valence character. The lifetimes of these states are quite different. The

Fig. 1 Experimental setup and XUV spectra. **a** Schematic of the experimental setup. A broad XUV spectrum is generated by focusing a 10 fs NIR pulse into a xenon gas jet to generate an attosecond pulse train. The XUV is refocused in a second gas jet containing the target gas. A longer NIR pulse is used to impulsively align the molecules in the second gas jet. **b** XUV spectra taken without the target gas (black curve), after absorption by N_2 at the delay times corresponding to alignment (blue curve) and anti-alignment (red curve).

Fig. 2 Experimental and calculated XUV transient absorption in aligned N₂. **a** Static absorption of isotropic N₂ molecules from 12.7 to 23.7 eV. The positions of vibrational levels of the valence $b^1\Pi_u$ and $b^1\Sigma_u^+$ states and of the Rydberg series built on $X^2\Sigma_g^+$, $A^2\Pi_u$ and $B^2\Sigma_u^+$ N₂⁺ cores are indicated, parallel and perpendicular transitions are labeled by red and blue lines respectively. **b** Experimental and **c** calculated differential absorption spectra around the full revival of aligned N₂ molecules. **d** Modulations of the spectrum at 15.28 eV (blue circle) and 17.14 eV (red circle). **e** Calculated alignment degree $\langle \cos^2 \theta \rangle - 1/3$.

Table 1 Assignment of N ₂ absorption spectrum.			
ϵ (eV)	Transition	Description	N ₂ ⁺ core
12.7–15.58	$(1\pi_u)(3\sigma_g) \rightarrow (1\pi_g)^2$	$b^1\Pi_u^a$	Valence
	$2\sigma_u \rightarrow 1\pi_g$		
	$3\sigma_g \rightarrow 3\sigma_u$	$b^1\Sigma_u^{+a}$	Valence
	$1\pi_u \rightarrow 1\pi_g$		
	$3\sigma_g \rightarrow n\pi\sigma_u$	CY ^b	$X^2\Sigma_g^+$
15.58–16.70	$3\sigma_g \rightarrow n\pi\pi_u$	WJ ^b	
	$1\pi_u \rightarrow 3s\sigma_g$	o_3^a	$A^2\Pi_u$
	$1\pi_u \rightarrow nd\delta_g$	WT ^b	
	$1\pi_u \rightarrow nd\sigma_g$	OT ^b	
	Unassigned	UN ^{c,d}	
16.70–18.75	$2\sigma_u \rightarrow nd\sigma_g$	HA ^{b,c}	$B^2\Sigma_u^+$
	$2\sigma_u \rightarrow nd\pi_g$	HE ^{b,c}	
	$2\sigma_u \rightarrow ns\sigma_g$	New OT ^{b,c}	

The experimental absorption spectrum of N₂, shown in Fig. 2a, consists of numerous overlapping vibrational series associated with states of Rydberg and valence character. The absorption features are summarized by the photon energy ϵ , the column labeled "Description" is the name of the electronic transitions in the "Transition" column, the "N₂⁺ core" column is the ionic state, to which the Rydberg series converge

CY Carroll Yoshino, WJ Worley Jenkins, WT Worley third, OT Ogawa Tanaka, UN Unassigned, HA Hopfield absorption, HE Hopfield emission

^aReference³⁰
^bReference³¹
^cReference³²
^dReference³³

states between 17 and 18.5 eV have short lifetimes of ~ 10 fs due to autoionization¹² and the absorption peaks show wide asymmetric Fano line shapes, whereas states below the ionization potential (15.58 eV) have long lifetime $> ps$ ^{28,29} and the absorption peaks show narrow symmetric Lorentzian line shapes. Following the assignment of^{30–33}, the absorption features are summarized in Table 1. We use red (blue) lines to label the excited states with $^1\Sigma_u^+$ ($^1\Pi_u$) symmetry. Due to the finite energy resolution of the XUV spectrometer (70 meV), the full structure of the excited states cannot be resolved. Figure 2b shows the differential absorption spectrum measured around the full rotational revival delay of N₂. The alignment-induced absorbance modulation can be seen across the whole energy range, which shows that both long-lived states and short-lived autoionizing states are sensitive to molecular alignment at the time of the

probe signal. Figure 2d shows lineouts versus time delay for the absorption peaks at 15.28 eV (marked by the blue arrow) and 17.14 eV (marked by the red arrow); the different modulation depth and polarity are evident.

Figure 2b clearly shows that there are two types of absorption features: those that maximize when the molecular axis is parallel to the XUV polarization direction ($\tau = 8.57$ ps), and those that maximize when the molecular axis is perpendicular ($\tau = 8.32$ ps). For example, at 15 eV, the direction changes, as it does at 18.2 eV. Visually one can identify regions of parallel or perpendicular electronic transitions, illustrating the power of ATAS with aligned molecules.

We now turn to O₂. Figure 3a shows the static absorption of isotropic O₂. The ground state may be written as $(1\sigma_g)^2(1\sigma_u)^2(2\sigma_g)^2(2\sigma_u)^2(3\sigma_g)^2(1\pi_u)^4(1\pi_g)^2 X^3\Sigma_g^-$. The adiabatic ionization energies corresponding to different final states of O₂⁺ are $(1\pi_g)^{-1} X^2\Pi_g = 12.07$ eV, $(1\pi_u)^{-1} A^4\Pi_u = 16.1$ eV, $(1\pi_u)^{-1} A^2\Pi_u = 17.05$ eV, $(3\sigma_g)^{-1} b^4\Sigma_g^- = 18.17$ eV, $(3\sigma_g)^{-1} B^2\Sigma_g^- = 20.3$ eV, $(2\sigma_u)^{-1} c^4\Sigma_u^- = 24.56$ eV^{34,37}, as shown by the thick gray lines in Fig. 3(a). Following the assignment of^{34–36}, the absorption features are summarized in Table 2. Similarly, we use red (blue) lines to label the excited states with $^3\Sigma_u^-$ ($^3\Pi_u$) symmetry. Figure 3b shows the differential absorption spectra and Fig. 3d shows lineouts for the absorption peak at 13.43 eV (marked by blue arrow) and 15.18 eV (marked by red arrow).

The experimental results in Figs. 2 and 3 show that the transient absorption signal indeed depends on the orientation of the molecules at the time of the XUV pulse, answering the question that we posed in the introduction. But it is not clear why this is true. The pump pulse creates a coherence between rotational states in the electronic ground state. At a later time the XUV pulse coherently excites the ensemble of molecules to a number of excited electronic states. The superposition of states results in an induced time-dependent polarization which will radiate an electric field for many picoseconds. Because the emission is out of phase with the XUV pulse, the time-integrated emission will appear as absorption^{5,9}. For an atom, the emission will decay exponentially over picoseconds or nanoseconds due to the lifetime of the excited state, and thereby determines the

Fig. 3 Experimental and calculated XUV transient absorption in aligned O_2 . **a** Static absorption of isotropic O_2 molecules from 12 to 22.9 eV. The positions of vibrational levels of Rydberg series built on several O_2^+ cores are indicated, parallel and perpendicular transition are labeled by red and blue lines respectively. **b** Experimental and **c** calculated differential absorption spectra around the full revival of aligned O_2 molecules. **d** Modulations of the spectrum at 13.43 eV (blue circle) and 15.18 eV (red circle). **e** Calculated alignment degree $\langle \cos^2 \theta \rangle - 1/3$.

Table 2 Assignment of O_2 absorption spectrum.

ϵ (eV)	Transition	Description	O_2^+ core
12.3–17.05	$1\pi_u \rightarrow 3s\sigma_g$	H^a	$a^4\Pi_u$
	$1\pi_u \rightarrow 4s\sigma_g$	$I^{b,c}$	
	$1\pi_u \rightarrow 3d\delta_g$	$I^{b,c}$	
	$1\pi_u \rightarrow 3d\sigma_g$	$I^{b,c}$	
	$1\pi_u \rightarrow 3s\sigma_g$	J^a	$A^2\Pi_u$
17.05–20.3	$3\sigma_g \rightarrow n\rho\sigma_u$	Strong ^a	$b^4\Sigma_g^-$
	$3\sigma_g \rightarrow n\rho\pi_u$	Weak ^a	
	$3\sigma_g \rightarrow n\rho\sigma_u$	Strong ^a	$B^2\Sigma_g^-$
	$3\sigma_g \rightarrow n\rho\pi_u$	Weak ^a	
	$2\sigma_u \rightarrow 3s\sigma_g$	Strong ^a	$c^4\Sigma_u^-$

The experimental absorption spectrum of O_2 , shown in Fig. 3a, consists of numerous overlapping vibrational series associated with states of Rydberg and valence character. The absorption features are summarized by the photon energy ϵ , the column labeled “Description” is the name of the electronic transitions in the “Transition” column, the “ O_2^+ core” column is the ionic state, to which the Rydberg series converge

^aReference³⁴

^bReference³⁵

^cReference³⁶

Fig. 4 Calculated induced dipole moment for a rigid rotor linear molecule based on N_2 . A rotational wavepacket is created in the ground state by the pump pulse, and the XUV pulse arrives at a later time t_x . If t_x is during a rotational revival (blue), then a greater polarization is induced. On the other hand, if t_x is when the molecules are perpendicular to the polarization, then fewer molecules are excited. A larger induced polarization signal causes a greater absorption in the frequency spectrum. The rotational constants are $B_0 = 2 \text{ cm}^{-1}$ and $B_1 = 1.9 \text{ cm}^{-1}$.

spectral linewidth of the absorption feature. For molecules, the emitted polarization is more complicated, because the molecules continue to rotate after the XUV pulse.

Calculation of induced polarization. To understand the time-domain picture of re-emission, we illustrate the time-varying polarization for a simplified rigid rotor modeled on the N_2 molecule. We consider rotational states with J in the range 0–10. We start with only $J = 0, M = 0$ and create a rotational wavepacket in the ground electronic state $|0\rangle$ by coherently populating the even J s via Raman transitions. This determines the set of a_j . The rotational wavepacket has a full revival at $t = 8.5 \text{ ps}$, just after an anti-revival at $t = 8.2 \text{ ps}$. At a delayed time t_x , the XUV pulse creates population in an excited electronic state $|1\rangle$ through dipole transitions. We consider a parallel transition, i.e. one which is allowed when the XUV polarization is parallel to the molecular axis. Rotational selection rules require that $\Delta J = \pm 1$, so a set of odd- J rotational states are populated with complex amplitudes b_j , creating a second rotational wavepacket in the excited electronic state.

The induced polarization is proportional to the time-dependent dipole moment of the molecule.

$$d(t) = \langle \psi^1(t) | r | \psi^0(t) \rangle + c.c. = \sum_{j,j'} a_j b_{j'}^* e^{iE_j(t-t_x)} e^{-iE_{j'}^0 t} d_{||} \langle j'M | \cos \theta | JM \rangle + c.c., \quad (4)$$

here E_j^0 and E_j^1 are the energies of the eigenstates, including both rotational and electronic parts, $d_{||}$ is the parallel component of the electronic transition dipole moment in the molecular frame. $d(t)$ will have complex behavior in time, especially if the two electronic states have different rotational constants. See the Supplementary Note 1 for more details. Figure 4 shows a calculation of the induced polarization for this simplified molecule. The polarization results in free induction decay from the excited states. The re-emission is out of phase with the incident attosecond pulse, resulting in absorption lines in the transmitted spectrum^{5,9,10}. A stronger polarization signal is associated with stronger absorption. The high frequency oscillations are mainly determined by the electronic transition frequency (14 eV in this calculation, not visible on this time scale). The low frequency oscillations are induced by the coherence

between different rotational states and different rotational rates between the ground and excited states. This dipole leads to an absorption structure around 14 eV, which consists of several peaks, the separations of which are related to the rotational energy ($\sim$ meV, which cannot be resolved by our XUV spectrometer). Probe times t_x at the rotational revival (blue) and the anti-revival (red) are shown. The orientation-dependent behavior is evident during the first picosecond. For aligned molecules, the polarization signal is greatest immediately after excitation. For anti-aligned molecules, the signal is at first small, then increases as the molecules move into alignment 300 fs later. The overall signal is greater when the probe pulse arrives during the molecular alignment. This agrees with our experimental result: parallel transition shows stronger absorption when the XUV pulse arrives during a time of maximum alignment to the polarization direction.

One can understand this behavior classically. If the probe pulse arrives during a rotational revival, then more excited state population is produced for a parallel transition, and the excited molecules will also be parallel to the polarization direction. But the molecules in the ground and excited states will rotate at different rates, meaning that the induced dipole will modulate at their difference frequency. This leads to the modulations seen in Fig. 4.

Frequency domain model. We now proceed to model the expected absorption in the frequency domain. More details are provided in the Supplementary Note 2. Impulsive alignment of a rigid rotor results in a distribution of molecular angles θ relative to the polarization axis $\hat{z}$ of the laser pulse. The distribution is cylindrically symmetric about $\hat{z}$ and varies in time based on the rotational constant of the molecule and its nuclear spin statistics¹⁹. The degree of alignment is characterized by the expectation value of the cosine-squared of the angle of the molecular axes to $\hat{z}$, namely $\langle \cos^2 \theta \rangle(t)$. For an isotropic sample, $\langle \cos^2 \theta \rangle = 1/3$; for a sample perfectly aligned along direction $\hat{z}$ it is 1. At characteristic times called rotational revivals, the molecular axis distribution achieves its maximum degree of alignment. Just before a full revival, the molecular axes lie mostly in a plane perpendicular to $\hat{z}$, call anti-alignment, quickly followed by a maximum degree of alignment parallel to $\hat{z}$. Figures 2e and 3e show the calculated evolution of $\langle \cos^2 \theta \rangle(t) - 1/3$ in the region of the rotational revival.

For molecules of $D_{\infty h}$ symmetry group, all bound-bound electronic transitions must be either parallel or perpendicular to the molecular axis³⁸. The measured absorption will then be a mixture of both types, weighted by the degree of alignment,

$$\sigma(E, t) = \langle \cos^2 \theta \rangle(t) \cdot \sigma_{\parallel}(E) + [1 - \langle \cos^2 \theta \rangle(t)] \cdot \sigma_{\perp}(E), \quad (5)$$

where $\sigma_{\parallel}(E)$ and $\sigma_{\perp}(E)$ are absorption cross sections for parallel and perpendicular transitions. The absorption cross section for bound-bound transitions are taken from the high-resolution spectra of ref. ²⁵ for N_2 and ref. ³⁴ for O_2 , including their symmetry assignments (summarized in Tables 1 and 2). The absorption cross sections for bound-free transitions above each ionization threshold are assumed to be flat in energy and are fit to the experiment, as described in the Supplementary Note 2. The calculated transmitted spectra are convolved with the experimental instrumental response function of 70 meV width, and the predicted differential absorption spectrum ΔOD is calculated in the same way as for the experimental data. The model results are shown in Fig. 2c for N_2 and Fig. 3c for O_2 .

The experimental and the calculated results are in good agreement. The significant difference around 13 eV for N_2 may come from several factors. First, the XUV signal itself is weak around 13 eV as shown in Fig. 1(b). Second, the absorption cross

section is huge ($\sim$ 400 Mb) as shown in the Supplementary Fig. 3, which means the XUV spectrum after absorption is very weak and the signal-to-noise ratio is poor in this energy range. Third, the second order diffraction of higher energy XUV photons (26 eV) falls onto 13 eV, which may pollute the signal. The simulated signal contrast is also larger than the measured contrast. One possible explanation is the assumption of full coherence of the rotational wavepacket in the ground state, by the use of Schrodinger's equation. A more accurate simulation would instead use a density matrix approach in which a degree of incoherence in the rotational wavepacket is included. Another possible reason is the assumption of a constant spectrometer resolution in the whole spectrum range in the simulation, whereas the real resolution changes with photon energy (better resolution at lower photon energy).

Discussion

The photoabsorption spectra of small molecules have been extensively studied using high-resolution XUV monochromators at synchrotron radiation facilities^{25,34,39–42}. Large numbers of observed bands in the absorption spectra have been grouped into Rydberg series converging to different vibrational levels of various ion states. But one cannot get information on symmetries of photoexcited states from traditional photoabsorption spectroscopy without the help of theoretical calculations⁴³. The Rydberg orbitals are usually determined by calculating the effective quantum number n^* and quantum defect δ , then the symmetry of the Rydberg state can be classified according to the build-up principles given by Herzberg⁴⁴. Normally, assignments can be confirmed from the rotational analysis of the Rydberg transition or from the analysis of fluorescence polarization. However, few such data are available. In O_2 , the lifetime broadening due to rapid autoionization and/or predissociation results in the absence of clearly resolved rotational structure^{34,45}. So the symmetry assigned to some Rydberg series are tentative and various interpretations exist^{35,36,46–50}.

Despite extensive experimental and theoretical work on the assignments of the Rydberg series converging onto the N_2^+ X, A, and B states, a consistent interpretation has yet to emerge⁴⁶. The main difference between these assignments concerns the Rydberg series converging onto the N_2^+ A threshold. In Fig. 5(a), the black line is the absorption spectrum adapted from a synchrotron study²⁵. Two series known as Worley's third series and Ogawa-Tanaka series have been observed, labeled as WT and OT. However, some of the most prominent features in the absorption spectrum are not included in either of these two series and are labeled as UN. Figure 5b is a magnified portion of Fig. 2b from 15.72 to 16.76 eV. We can see most of the UN peaks are consistent with parallel transitions (negative for anti-alignment and positive for alignment).

For better comparison, we choose the alignment and anti-alignment delay points, labeled by the red and blue arrows in Fig. 5b. We define an asymmetry signal: $P = \Delta OD(\text{Alignment}) - \Delta OD(\text{Anti alignment})$, shown by the black line in Fig. 5c. If we assume that the UN series have $^1\Sigma_u^+$ symmetry (parallel transitions), the predicted spectrum is shown by the red line. On the other hand, assuming $^1\Pi_u$ symmetry (perpendicular transitions) gives the blue line. The red line shows much better agreement with the experimental result, which suggests that these unassigned peaks have $^1\Sigma_u^+$ symmetry. Consider the Rydberg orbital to be $n l \lambda_{g/u}$, where n is the principal quantum number, l is the orbital angular momentum quantum number, λ is the quantum number of the projection of orbital angular momentum onto the internuclear axis and g/u means symmetric/antisymmetric orbital. Since the vibrational spacings in the UN series resemble those of the $N_2^+ A^2\Pi_u$ state^{32,33}, these series should originate from the

Fig. 5 Assigning symmetry to some Rydberg series of N₂ and O₂. **a** High resolution absorption spectrum of N₂ adapted from²⁵, UN refers to unassigned states. **b** Zoom-in of Fig. 2b. **c** Measured and calculated asymmetry signals. **d** High resolution absorption spectrum of O₂ adapted from ref. ³⁴. **e** Magnification of Fig. 3b. **f** Measured and calculated asymmetry signals.

promotion of a $1\pi_u$ electron. So the Rydberg orbital symmetry must be g , which means l is even⁴⁴. Our experiment identified that the UN series are parallel transitions, so λ should be π , which means $\lambda = 1$. Since $l \geq \lambda$, l must be 2, which means a d orbital. We also calculated the quantum defects δ by $\nu_n(\nu) = \nu_\infty(\nu) - R/(n - \delta)^2$ ⁵⁰, where $\nu_n(\nu)$ is the Rydberg state with principal quantum number n and in the corresponding ν vibrational level³³, $\nu_\infty(\nu)$ is the series limit⁵¹, $R = 109737.316 \text{ cm}^{-1}$ is the Rydberg constant. For molecules built up from atoms of the first two rows of the periodic table, $\delta \sim 1$ for s orbital, $\delta \sim 0.7$ for p orbital and $\delta \sim 0$ for d orbital⁵². The result $\delta \sim -0.2$ also suggests a d orbital. So the transitions for these UN series are $1\pi_u \rightarrow nd\pi_g$. Usually quantum defect calculations are not enough for the Rydberg orbital assignment, in this case, $\delta \sim -0.2$ suggests a d orbital, but it can be $d\sigma_g$, $d\pi_g$, or $d\delta_g$. Sometimes, even δ value itself is difficult to get, that is why the questions about orbital assignment still exist even for simple diatomic molecules. Our assignment illustrates the value of transient absorption measurements made in the molecular frame.

For O₂, disputed structures exist around 15 eV. The adapted absorption spectrum from³⁴ is shown by the black line in Fig. 5d. Three progressions labeled as I, I' and I'' were observed, and converge onto the O₂⁺ $a^4\Pi_u$ threshold. The Rydberg orbital assignment of these states has been the subject of considerable debate^{35,36,49,50}. Similarly, Fig. 5e shows a magnified portion of Fig. 3b. Although the bound-free transition induces a parallel modulation background, we can still see that most I peaks fall onto the weak modulation part while most I' peaks fall onto the strong modulation. The black line in Fig. 5(f) shows the asymmetry signal. By assuming $^3\Pi_u$ and $^3\Sigma_u^-$ symmetries for I and I' respectively, the calculation shown by the red line in Fig. 5f achieved the best agreement. The effective quantum number of I' is $n^* = 2.9914$ ⁵⁰, indicating the corresponding transition is $1\pi_u \rightarrow 3d\pi_g$. The $^3\Sigma_u^-$ symmetry assignment for I' conflicts with most previous studies. A recent photodissociation study of O₂ around 14.6–15.2 eV also discovered some $^3\Sigma_u^-$ features around vibrational states ($\nu = 4, \nu = 5$) of I'⁵³. However, the most prominent features of I' is $\nu = 6$ and $\nu = 7$ ³⁴, which are beyond the reported spectral range of ref. ⁵³. The I'' state has much smaller absorption cross section and strongly overlaps with I and I', which makes the symmetry identification difficult. In the future, combining more highly aligned molecular ensembles and a higher

XUV spectral resolution, it shall be possible to resolve these features.

In conclusion, XUV transient absorption spectroscopy was demonstrated with aligned molecules, even though the Rydberg states have long lifetimes compared to rotational timescales^{28,29}. The excitation step in the molecular frame is fast, and the dependence on molecular angle is not lost despite the complex modulation of the resulting free induction decay signal. We identified the symmetry of an unassigned series in N₂, and found a conflict with previous assignment of the I' series in O₂. Looking forward, further control could be obtained by adding another NIR pulse after the XUV pulse, to induce dipole coupling between the excited electronic states, which can be controlled by alignment of the molecular target⁵⁴, allowing for a clearer physical interpretation of the complex absorption spectrum. We demonstrated XUV transient absorption spectroscopy in the molecular frame using linear molecules due to their simplicity in alignment. More complex molecules can be aligned in three dimensions by using a pulse sequence²⁴, permitting similar spectroscopic studies to be applied to symmetric top molecules.

Methods

Experimental setup. In the experiment, NIR laser pulses were generated by a Ti:sapphire laser system (100 Hz, 14 mJ, 800 nm, 50 fs). Approximately 1.6 mJ was sent into a differentially pumped hollow core fiber (250 μm inner diameter) filled with 0.7 bar of argon to create a 10 fs pulse centered around 800 nm. This pulse was focused into a pulsed gas jet of xenon (3 bar backing pressure, 250 μm nozzle) where high harmonic generation took place. A thin ($80 \pm 10 \mu\text{m}$) monocrystalline quartz plate was placed $\sim 2 \text{ cm}$ before the xenon gas jet to broaden the spectrum through self-phase modulation (SPM) and to create some second harmonic of 800 nm^{55,56}. The intensity at the quartz plate was $3 \times 10^{13} \text{ W/cm}^2$, below the damage threshold ($> 4 \times 10^{13} \text{ W/cm}^2$) but strong enough to generate new spectral components through SPM. The bandwidth of each harmonic increased from 0.7 to 2.3 eV, the total spectrum spanning 12–24 eV. The XUV was composed of a train of about five pulses in an attosecond pulse train. A 500- μm -diameter pinhole was used to block the residual NIR driving field and transmit most of the XUV radiation. An intensity of $2 \times 10^{11} \text{ W/cm}^2$ is estimated for the residual NIR beam at the sample target based on the Stark shift (see Supplementary Note 3 and Supplementary Fig. 4). All the results in this article are reproducible when we replace the pinhole with a 100-nm-thick aluminum foil (transparent energy range: 15–73 eV) or a 350-nm-thick indium foil (transparent energy range: 11–17 eV).

A part (1 mJ) of the initial 50 fs NIR pulse was combined with the XUV pulse with a holey mirror. Both the XUV and the NIR pulses propagated noncollinearly and intersected at the absorption gas jet (250 μm nozzle, gas density $\sim 10^{18} \text{ cm}^{-3}$, interaction length $\sim 0.5 \text{ mm}$; N₂: 99.999% purity, 10 bar backing pressure, O₂:

99.999% purity, 6.5 bar backing pressure) with a crossing angle of 18 mrad. The NIR pulse at the target (3×10^{13} W/cm², pulse duration increased to 70 fs due to the dispersion of transmission optics) was used to perform impulsive alignment on the sample molecules. The transmitted XUV spectrum was dispersed in a flat-field XUV spectrometer and detected by a microchannel plate detector, with a spectral resolution of 70 meV.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Code availability

The codes used in this study are available from the corresponding author upon reasonable request.

Received: 3 June 2019; Accepted: 29 October 2019;

Published online: 21 November 2019

References

- Goulielmakis, E. et al. Real-time observation of valence electron motion. *Nature* **466**, 739–743 (2010).
- Ott, C. et al. Lorentz meets fano in spectral line shapes: a universal phase and its laser control. *Science* **340**, 716–720 (2013).
- Ott, C. et al. Reconstruction and control of a time-dependent two-electron wave packet. *Nature* **516**, 374–378 (2014).
- Chini, M. et al. Sub-cycle oscillations in virtual states brought to light. *Sci. Rep.* **3**, 1105 (2013).
- Beck, A. R., Neumark, D. M. & Leone, S. R. Probing ultrafast dynamics with attosecond transient absorption. *Chem. Phys. Lett.* **624**, 119–130 (2015).
- Cao, W., Warrick, E. R., Fidler, A., Leone, S. R. & Neumark, D. M. Excited-state vibronic wave-packet dynamics in H₂ probed by xuv transient four-wave mixing. *Phys. Rev. A* **97**, 023401 (2018).
- Warrick, E. R., Cao, W., Neumark, D. M. & Leone, S. R. Probing the dynamics of rydberg and valence states of molecular nitrogen with attosecond transient absorption spectroscopy. *J. Phys. Chem. A* **120**, 3165–3174 (2016).
- Warrick, E. R. et al. Attosecond transient absorption spectroscopy of molecular nitrogen: vibrational coherences in the b' ¹Σ_u⁺ state. *Chem. Phys. Lett.* **683**, 408–415 (2017).
- Wu, M., Chen, S., Camp, S., Schafer, K. J. & Gaarde, M. B. Theory of strong-field attosecond transient absorption. *J. Phys. B: At. Mol. Opt. Phys.* **49**, 062003 (2016).
- Santra, R., Yakovlev, V. S., Pfeifer, T. & Loh, Z.-H. Theory of attosecond transient absorption spectroscopy of strong-field-generated ions. *Phys. Rev. A* **83**, 033405 (2011).
- Cheng, Y. et al. Reconstruction of an excited-state molecular wave packet with attosecond transient absorption spectroscopy. *Phys. Rev. A* **94**, 023403 (2016).
- Reduzzi, M. et al. Observation of autoionization dynamics and sub-cycle quantum beating in electronic molecular wave packets. *J. Phys. B: At. Mol. Opt. Phys.* **49**, 065102 (2016).
- Leone, S. R. et al. What will it take to observe processes in 'real time'? *Nat. Photon.* **8**, 162–166 (2014).
- Hosler, E. R. & Leone, S. R. Characterization of vibrational wave packets by core-level high-harmonic transient absorption spectroscopy. *Phys. Rev. A* **88**, 023420 (2013).
- Pertot, Y. et al. Time-resolved x-ray absorption spectroscopy with a water window high-harmonic source. *Science* **355**, 264–267 (2017).
- Attar, A. R. et al. Femtosecond x-ray spectroscopy of an electrocyclic ring-opening reaction. *Science* **356**, 54–59 (2017).
- Popmintchev, D. et al. Near- and extended-edge x-ray-absorption fine-structure spectroscopy using ultrafast coherent high-order harmonic supercontinua. *Phys. Rev. Lett.* **120**, 093002 (2018).
- Kobayashi, Y., Chang, K. F., Zeng, T., Neumark, D. M. & Leone, S. R. Direct mapping of curve-crossing dynamics in ibr by attosecond transient absorption spectroscopy. *Science* **365**, 79–83 (2019).
- Stapelfeldt, H. & Seideman, T. Colloquium: aligning molecules with strong laser pulses. *Rev. Mod. Phys.* **75**, 543–557 (2003).
- Peng, P., Bai, Y., Li, N. & Liu, P. Measurement of field-free molecular alignment by balanced weak field polarization technique. *AIP Adv.* **5**, 127205 (2015).
- Thomann, I. et al. Direct measurement of the angular dependence of the single-photon ionization of aligned N₂ and CO₂. *J. Phys. Chem. A* **112**, 9382–9386 (2008).
- Li, J. et al. 53-attosecond x-ray pulses reach the carbon k-edge. *Nat. Commun.* **8**, 186 (2017).
- Gaumnitz, T. et al. Streaking of 43-attosecond soft-x-ray pulses generated by a passively cep-stable mid-infrared driver. *Opt. Express* **25**, 27506–27518 (2017).
- Lee, K. F., Villeneuve, D. M., Corkum, P. B., Stolow, A. & Underwood, J. G. Field-free three-dimensional alignment of polyatomic molecules. *Phys. Rev. Lett.* **97**, 173001 (2006).
- Gürtler, P., Saile, V. & Koch, E. High resolution absorption spectrum of nitrogen in the vacuum ultraviolet. *Chem. Phys. Lett.* **48**, 245–250 (1977).
- Lofthus, A. & Krupenie, P. H. The spectrum of molecular nitrogen. *J. Phys. Chem. Ref. Data* **6**, 113–307 (1977).
- Yencha, A. J., Ellis, K. & King, G. C. High-resolution threshold photoelectron and photoion spectroscopy of molecular nitrogen in the 15.0–52.7 eV photon energy range. *J. Electron Spectrosc. Relat. Phenom.* **195**, 160–173 (2014).
- Oertel, H., Kratzat, M., Imschweiler, J. & Noll, T. Fluorescence from ¹Π_u and ¹Σ_u⁺ states of molecular nitrogen excited with synchrotron radiation between 12.4 and 18.8 eV. *Chem. Phys. Lett.* **82**, 552–556 (1981).
- Sprengers, J. et al. Pump–probe lifetime measurements on singlet ungerade states in molecular nitrogen. *Chem. Phys. Lett.* **389**, 348–351 (2004).
- Pratt, S. T., Dehmer, P. M. & Dehmer, J. L. Photoelectron studies of resonant multiphoton ionization of molecular nitrogen. *J. Chem. Phys.* **81**, 3444–3451 (1984).
- Kosman, W. M. & Wallace, S. Complete dipole oscillator strength distribution and its moments for N₂. *J. Chem. Phys.* **82**, 1385–1399 (1985).
- Berg, L.-E., Erman, P., Källne, E., Sorensen, S. & Sundström, G. Studies of photoionization and photodissociation of N₂ in the 15–30 eV region using intense synchrotron radiation. *Phys. Scr.* **44**, 131–137 (1991).
- Ogawa, M. Vibrational isotope shifts of absorption bands of n2 in the spectral region 720–830 Å. *Can. J. Phys.* **42**, 1087–1096 (1964).
- Holland, D. et al. A study of the absolute photoabsorption, photoionization and photodissociation cross sections and the photoionization quantum efficiency of oxygen from the ionization threshold to 490 Å. *Chem. Phys.* **173**, 315–331 (1993).
- Cubric, D., Wills, A. A., Comer, J. & Ukai, M. Selective population of spin-orbit levels in the autoionization of O₂. *J. Phys. B: At. Mol. Opt. Phys.* **26**, 3081–3099 (1993).
- Demekhin, P. V., Sukhorukov, V. L., Schmoranz, H. & Ehresmann, A. Neutral dissociation of the I, I', and I'' vibronic progressions of O₂. *J. Chem. Phys.* **132**, 204303 (2010).
- Krupenie, P. H. The spectrum of molecular oxygen. *J. Phys. Chem. Ref. Data* **1**, 423–534 (1972).
- Atkins, P. W. & Friedman, R. S. *Molecular Quantum Mechanics* (Oxford University Press, 2011).
- Shaw, D. et al. A study of the absolute photoabsorption, photoionisation and photodissociation cross sections and the photoionisation quantum efficiency of carbon dioxide from the ionisation threshold to 345 Å. *Chem. Phys.* **198**, 381–396 (1995).
- Shaw, D. et al. A study of the absolute photoabsorption cross section and the photoionization quantum efficiency of nitrous oxide from the ionization threshold to 480 Å. *Chem. Phys.* **163**, 387–404 (1992).
- Berkowitz, J. *Atomic and Molecular Photoabsorption: Absolute Total Cross Sections* (Academic Press, 2002).
- Berkowitz, J. *Atomic and Molecular Photoabsorption: Absolute Partial Cross Sections* (Academic Press, 2015).
- ichi Adachi, J., Kosugi, N. & Yagishita, A. Symmetry-resolved soft x-ray absorption spectroscopy: its application to simple molecules. *J. Phys. B: At. Mol. Opt. Phys.* **38**, R127–R152 (2005).
- Herzberg, G. *Molecular Spectra and Molecular Structure. Vol. 1: Spectra of Diatomic Molecules* (Van Nostrand Reinhold Co., New York, 1950).
- Tonkyn, R. G., Winniczek, J. W. & White, M. G. Photoionization and photoelectron spectroscopy of o2 with coherent vacuum ultraviolet radiation. *J. Chem. Phys.* **91**, 6632–6642 (1989).
- Shaw, D. et al. A study of the absolute photoabsorption cross section and the photoionization quantum efficiency of nitrogen from the ionization threshold to 485 Å. *Chem. Phys.* **166**, 379–391 (1992).
- Carter, V. L. High-resolution n2 absorption study from 730 to 980 Å. *J. Chem. Phys.* **56**, 4195–4205 (1972).
- Dehmer, P. M. & Chupka, W. A. High resolution study of photoionization processes in O₂. *J. Chem. Phys.* **62**, 4525–4534 (1975).
- Katayama, D. & Tanaka, Y. The I, I', and I'' absorption bands of O₂ in the region 790–865 Å. *J. Mol. Spectrosc.* **88**, 41–50 (1981).
- Wu, C. R. Assignments of autoionization states of O₂. *J. Quantitative Spectrosc. Radiative Transfer* **37**, 1–15 (1987).
- Yoshino, K., Ogawa, M. & Tanaka, Y. Extension of rydberg absorption series of N₂, A²Π_u ← X¹Σ_g⁺. *J. Mol. Spectrosc.* **61**, 403–411 (1976).
- Lefebvre-Brion, H. & Field, R. W. *The Spectra and Dynamics of Diatomic Molecules: Revised and Enlarged Edition* (Elsevier, 2004).
- Zhou, Y., Meng, Q. & Mo, Y. Photodissociation dynamics of superexcited O₂: dissociation channels O(⁵S) vs. O(³S). *J. Chem. Phys.* **141**, 014301 (2014).
- Bækhoj, J. E. & Madsen, L. B. Attosecond transient-absorption spectroscopy on aligned molecules. *Phys. Rev. A* **94**, 043414 (2016).

55. Hammond, T. J. et al. Integrating solids and gases for attosecond pulse generation. *Nat. Photonics* **11**, 594–599 (2017).
56. Hammond, T. J., Villeneuve, D. M. & Corkum, P. B. Producing and controlling half-cycle near-infrared electric-field transients. *Optica* **4**, 826–830 (2017).

Acknowledgements

The authors thank D. Crane and R. Kroeker for technical support. The authors acknowledge fruitful discussions with X. Ding, T. J. Hammond, M. S. Schuurman, and M. Spanner. This research is supported by the NSERC Discovery Grant program. In addition, we acknowledge the support of the Joint Centre for Extreme Photonics.

Author contributions

P.P. and D.M.V. conceived and planned the experiment. P.P., C.M., and M.H. conducted the measurements. P.P., C.M., and D.M.V. analyzed and interpreted the data. P.B.C., A.Y. N., and D.M.V. supervised the project. All authors discussed the results and contributed to the final manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41467-019-13251-w>.

Correspondence and requests for materials should be addressed to P.P. or D.M.V.

Peer review information *Nature Communications* thanks the anonymous reviewers for their contributions to the peer review of this work.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019

Supplementary Information

Symmetry of Molecular Rydberg States Revealed by XUV Transient Absorption Spectroscopy

Peng *et al.*

Supplementary Note 1: Time Response of Induced Polarization

The main text showed the time-dependent response of aligned molecules probed at different times. We approximate the N_2 molecule as a rigid rotor with a rotational constant $B_0 = 2 \text{ cm}^{-1}$. With an initial rotational temperature of $T = 0$, only the $J = 0, M = 0$ rotational state is populated. We simulate the alignment process by equally populating the even J states, $J = 0, 2, 4, 6, 8, 10$, since the Raman process requires $\Delta J = \pm 2$. The degree of alignment is shown in Supplementary Fig. 1. The full revival is around $t = 8.5 \text{ ps}$ at which time the molecular axes are maximally parallel to the polarization axis. At $t = 8.2 \text{ ps}$, the molecules are maximally perpendicular to the polarization axis (this simplified model can not reproduce 1/4 and 3/4 revivals accurately, however, it is enough for simulating full revival and is beneficial to give a physical insight to the time response of induced polarization).

The ground state wavefunction is written as

$$|\psi^0\rangle = \sum_J a_J e^{-iE_J^0 t} |0\rangle |JM\rangle, \quad (1)$$

where $E_J^0 = B_0 J(J+1)$ is the rotational energy. At a delayed time t_x , a delta-function pulse creates population in the excited electronic state,

$$|\psi^1\rangle = \sum_J b_J e^{-iE_J^1(t-t_x)} |1\rangle |JM\rangle, \quad (2)$$

where $E_J^1 = E_1 + B_1 J(J+1)$ is the energy of the excited rotational states, including the electronic term energy E_1 . The values of b_J are determined by time-dependent perturbation theory. The induced dipole moment is

$$\begin{aligned} d(t) &= \langle \psi^1(t) | r | \psi^0(t) \rangle + c.c. \\ &= \sum_{JJ'} a_J b_{J'}^* e^{iE_{J'}^1(t-t_x)} e^{-iE_J^0 t} \times d_{\parallel} \langle J'M | \cos \theta | JM \rangle + c.c., \end{aligned} \quad (3)$$

where $d_{\parallel}$ is the electronic transition dipole moment. It is $d(t)$ that is plotted in Fig. 4 of the main text.

Fig. 4 of the main text shows the induced polarization for the case when the rotational constant of the upper and lower states are different, which is typical of almost all systems. In Supplementary Fig. 2 we show a calculation when $B_0 = B_1 = 2 \text{ cm}^{-1}$. Because both the ground and excited state molecules are "rotating" at the same speed, the signal appears less chaotic. The periodicity of the ground state alignment (Supplementary Fig. 1) is more apparent. Both Supplementary Fig. 2 and Fig. 4 of the main text show the polarization signal is greatest immediately after excitation for aligned molecules, whereas for anti-aligned molecules, the signal is at first small, then increases as the molecules move into alignment 300 fs later. The overall signal is greater when the probe pulse arrives during the molecular alignment. This agrees with our experimental result: parallel transition shows stronger absorption when the XUV pulse arrives during a time of maximum alignment to the polarization direction.

Supplementary Fig. 1. Calculated degree of alignment of the simulated N_2 molecules. The XUV probe pulse arrives when the molecules are maximally aligned parallel or perpendicular to the polarization axis, as indicated by the arrows.

Supplementary Fig. 2. Calculated induced dipole moment for a rigid rotor linear molecule based on N_2 . A rotational wavepacket is created in the ground state by the pump pulse, and the XUV pulse arrives at a later time t_x . If t_x is during a rotational revival (blue), then a greater polarization is induced. On the other hand, if t_x is when the molecules are perpendicular to the polarization, then fewer molecules are excited. The rotational constants are $B_0 = 2 \text{ cm}^{-1}$ and $B_1 = 2 \text{ cm}^{-1}$.

Supplementary Note 2: Frequency Domain Model

Molecular Alignment

When linear molecules are irradiated by a pump pulse whose duration $\tau = 70$ fs is much shorter than the rotational period, nonadiabatic field-free alignment is achieved [1, 2]. Each initial rotational eigenstate $|J_0, M_0\rangle$ will expand to a rotational wave packet $\Psi_{J_0, M_0}(t) = \sum_{J, M} a_{J, M}^{J_0, M_0}(t) |J, M\rangle$, where J is the rotational quantum number, M is the projection on the polarization axis and $a_{J, M}^{J_0, M_0}(t)$ is the amplitude for each $|J, M\rangle$ state. The time evolution of the wave packet can be calculated by solving the time-dependent Schrodinger equation. The degree of alignment is characterized by $\langle \cos^2 \theta \rangle_{J_0, M_0}(t) = \langle \Psi_{J_0, M_0}(t) | \cos^2 \theta | \Psi_{J_0, M_0}(t) \rangle$, in which θ is the angle of the molecular axis to the polarization axis. Finally, thermally averaged values of the degree of alignment are obtained as:

$$\langle \cos^2 \theta \rangle(t) = \sum_{J_0, M_0} \rho_{J_0, M_0}(T) \langle \cos^2 \theta \rangle_{J_0, M_0}(t) \quad (4)$$

where T is the rotational temperature and $\rho_{J_0, M_0}(T)$ is the Boltzmann weight function for the $|J_0, M_0\rangle$ initial state. To calculate the rotational wavepacket created by the pump pulse, we use $I = 3 \times 10^{13}$ W/cm², $\tau = 70$ fs and a rotational temperature of 100 K. The calculated $\langle \cos^2 \theta \rangle(t) - 1/3$ is shown in Fig. 2(e) and Fig. 3(e) of the main manuscript.

Absorption Cross Section

The absorption cross section contains contributions from both parallel and perpendicular transitions in the molecular frame, depending on the degree of alignment:

$$\sigma(E, t) = \langle \cos^2 \theta \rangle(t) \cdot \sigma_{\parallel}(E) + [1 - \langle \cos^2 \theta \rangle(t)] \cdot \sigma_{\perp}(E), \quad (5)$$

where $\sigma_{\parallel}(E)$ and $\sigma_{\perp}(E)$ are absorption cross sections for parallel and perpendicular contributions, $\langle \cos^2 \theta \rangle(t)$ is the alignment degree. For an isotropic medium, $\langle \cos^2 \theta \rangle(t) = \frac{1}{3}$, and so $\sigma(E) = \frac{\sigma_{\parallel}(E)}{3} + \frac{2\sigma_{\perp}(E)}{3}$ [3].

Bound-Bound and Bound-Free Transitions

We consider two types of dipole-allowed transitions initiated by the XUV: bound-bound and bound-free. As N₂ and O₂ are linear molecules of D_{∞h} symmetry, with the help of the direct product table of Supplementary reference [4], all the bound-bound transitions are pure parallel or pure perpendicular. As our XUV spectrum covers several ionization thresholds of N₂ and O₂, bound-free transitions will induce a continuum absorption background. These bound-free transitions can be of dominant parallel or perpendicular character. The total absorption cross section at different alignments is defined as:

$$\begin{aligned} \sigma_{\text{total}}(E, t) = & \Sigma_m \langle \cos^2 \theta \rangle(t) \cdot \sigma_{m\parallel}(E) + \Sigma_n [1 - \langle \cos^2 \theta \rangle(t)] \cdot \sigma_{n\perp}(E) \\ & + \Sigma_c [\langle \cos^2 \theta \rangle(t) \cdot \sigma_{c\parallel}(E) + [1 - \langle \cos^2 \theta \rangle(t)] \cdot \sigma_{c\perp}(E)]. \end{aligned} \quad (6)$$

The three terms in the summation correspond to bound-bound parallel, bound-bound perpendicular and bound-free transitions, respectively.

Supplementary Fig. 3. Absorption cross section of N_2 . (a) High resolution absorption spectrum of N_2 adapted from [5]. Black line: total bound-bound and bound-free absorption cross section. Magenta line: total bound-free absorption cross section. (b) Partial bound-free absorption cross section.

N_2 Example

For N_2 , the black line in Supplementary Fig. 3(a) shows the high resolution absorption spectrum adapted from [5]. We get the values of $\sigma_{m\parallel}(E)$ and $\sigma_{n\perp}(E)$ from the discrete part of the absorption cross section. Modelling the bound-free part is a bit more complicated. $\sigma_{c\parallel}(E)$ and $\sigma_{c\perp}(E)$ are bound-free absorption cross sections for parallel and perpendicular contribution, c can be HOMO, HOMO-1 and HOMO-2 orbitals in our situation. The continuum baseline shown by the magenta line Supplementary Fig. 3(a) gives the total bound-free absorption cross section. Combining with the ionization ratios for different orbitals [6, 7], we can get the partial bound-free absorption cross section: $\sigma_{\text{HOMO}}(E)$, $\sigma_{\text{HOMO-1}}(E)$ and $\sigma_{\text{HOMO-2}}(E)$, shown in Supplementary Fig. 3(b). But $\sigma_{c\parallel}(E)$ and $\sigma_{c\perp}(E)$ for each orbital are still unknown. In principle one can calculate $\sigma_{c\parallel}(E)$ and $\sigma_{c\perp}(E)$, but there are differences according to the computational method used [8–11]. As these bound-free absorption cross sections usually vary slowly with photon energy, we define a constant ratio $\alpha_c = \sigma_{c\parallel}(E)/\sigma_c(E)$. From $\sigma_c(E) = \frac{\sigma_{c\parallel}(E)}{3} + \frac{2\sigma_{c\perp}(E)}{3}$, we know $0 \leq \alpha_c \leq 3$, this value may vary with orbitals. Therefore, there are only three unknown constants left in Supplementary Eq. (6), which are α_{HOMO} , $\alpha_{\text{HOMO-1}}$ and $\alpha_{\text{HOMO-2}}$.

Beer-Lambert Law and Resolution Considerations

The XUV spectrum after it interacts with the macroscopic gas sample was calculated by using Beer-Lambert law:

$$I(E, t) = I_0(E) e^{-\rho \sigma_{\text{total}}(E, t) z}, \quad (7)$$

where $I_0(E)$ is the measured XUV spectrum without absorption gas shown by the black line in Fig. 1(b) of

the main manuscript, ρ is the gas density, z is the medium length, $\sigma_{\text{total}}(E, t)$ is the total absorption cross section shown in Supplementary Eq. (6). We determine the gas density by using the smooth continuum part of the measured static absorption of isotropic molecules, and infer $\rho(\text{N}_2) = 1.177 \times 10^{18} \text{ cm}^{-3}$ and $\rho(\text{O}_2) = 9.77 \times 10^{17} \text{ cm}^{-3}$ for a medium length of 0.5 mm. The calculated transmitted spectrum is convolved with the experimental spectral resolution to get:

$$I_c(E, t) = I(E, t) * G(E), \quad (8)$$

where $G(E) = e^{-4 \ln(2) (\frac{E}{\delta E})^2}$ and $\delta E = 70 \text{ meV}$.

Differential Absorption Spectrum

For isotropic sample, the total absorption cross section was given by the high resolution synchrotron data, shown by the black line in Supplementary Fig. 3(a) for N_2 . Combining with Supplementary Eq. (7) and Supplementary Eq. (8), we can get the XUV spectrum after transmission through the isotropic sample $I_{\text{off}}(E)$.

For aligned sample, the total absorption cross section was calculated by Supplementary Eq. (6), and we can calculate the XUV spectrum after transmission through the aligned sample $I_{\text{on}}(E, t)$.

The predicted differential absorption spectrum is calculated in the same way as for the experimental data:

$$\Delta\text{OD}(E, t) = -\log_{10}[I_{\text{on}}(E, t)/I_{\text{off}}(E)]. \quad (9)$$

Comparing with the measured ΔOD , α_{HOMO} , $\alpha_{\text{HOMO}-1}$ and $\alpha_{\text{HOMO}-2}$ were determined by a fitting procedure. For N_2 , $\alpha_{\text{HOMO}} = 2.1$, $\alpha_{\text{HOMO}-1} = 0.165$, $\alpha_{\text{HOMO}-2} = 1.6$. For O_2 , $\alpha_{\text{HOMO}} = 1.3$, $\alpha_{\text{HOMO}-1} = 0.6$, $\alpha_{\text{HOMO}-2} = 2.5$. The calculation results were shown in Fig. 2(c) and Fig. 3(c) of the main manuscript. For both molecules, α values show parallel transition contribute more for HOMO and HOMO-2 orbitals while perpendicular transition contribute more for HOMO-1 orbital, which agrees with previous photonionization results [12–15].

Supplementary Note 3: Residual NIR Driving Pulse Intensity

We estimated the NIR intensity by comparing the static absorption of N_2 measured with an indium filter and with a pinhole (the NIR alignment pulse was blocked in both measurements). As shown in Supplementary Fig. 4, the absorption spectrum were normalized by the peak nearest to the ionization potential (15.58 eV). The residual NIR driving field will shift the absorption peak to higher energy due to Stark effect (the relative peak heights are also different in these two measurements, because the XUV spectrum shape will change after passing through the In filter). The shift amount is ~ 12 meV, from : $U_p = e^2 E^2 / 4mw^2$ and $I = c\varepsilon_0 E^2 / 2$, where U_p is the ponderomotive energy, e is the electronic charge, E is the electric field, m is the mass of the electron, w is the pulse frequency, I is the pulse intensity, ε_0 is the permittivity of vacuum. The estimated residual NIR intensity is 2×10^{11} W/cm². The absorption linewidths in two measurements are similar which also indicate the residual NIR pulse is weak.

Supplementary Fig. 4. Absorption spectrum of N_2 around ionization potential. Red line: measured with a pinhole, blue line: measured with an indium filter.

Supplementary References

1. Stapelfeldt, H. & Seideman, T. Colloquium: Aligning molecules with strong laser pulses. *Rev. Mod. Phys.* **75**, 543–557 (2003).
2. Peng, P., Bai, Y., Li, N. & Liu, P. Measurement of field-free molecular alignment by balanced weak field polarization technique. *AIP Advances* **5**, 127205 (2015).
3. Cheng, Y. *et al.* Reconstruction of an excited-state molecular wave packet with attosecond transient absorption spectroscopy. *Phys. Rev. A* **94**, 023403 (2016).
4. Atkins, P. W. & Friedman, R. S. *Molecular quantum mechanics* (Oxford university press, 2011).
5. Gürtler, P., Saile, V. & Koch, E. High resolution absorption spectrum of nitrogen in the vacuum ultraviolet. *Chemical Physics Letters* **48**, 245–250 (1977).
6. Plummer, E. W., Gustafsson, T., Gudat, W. & Eastman, D. E. Partial photoionization cross sections of N₂ and CO using synchrotron radiation. *Phys. Rev. A* **15**, 2339–2355 (1977).
7. Samson, J. A., Gardner, J. & Haddad, G. Total and partial photoionization cross-sections of O₂ from 100 to 800 Å. *Journal of Electron Spectroscopy and Related Phenomena* **12**, 281–292 (1977).
8. Kosman, W. M. & Wallace, S. Complete dipole oscillator strength distribution and its moments for N₂. *The Journal of Chemical Physics* **82**, 1385–1399 (1985).
9. Semenov, S. K., Cherepkov, N. A., Fecher, G. H. & Schönhense, G. Generalization of the atomic random-phase-approximation method for diatomic molecules: N₂ photoionization cross-section calculations. *Phys. Rev. A* **61**, 032704 (2000).
10. Lucchese, R. R., Takatsuka, K. & McKoy, V. Applications of the Schwinger variational principle to electron-molecule collisions and molecular photoionization. *Physics Reports* **131**, 147–221 (1986).
11. Gerwer, A., Asaro, C., McKoy, B. V. & Langhoff, P. W. Photoexcitation and ionization in molecular oxygen: Theoretical studies of electronic transitions in the discrete and continuous spectral intervals. *The Journal of Chemical Physics* **72**, 713–727 (1980).
12. Rouzée, A. *et al.* Photoelectron kinetic and angular distributions for the ionization of aligned molecules using a HHG source. *Journal of Physics B: Atomic, Molecular and Optical Physics* **45**, 074016 (2012).
13. Rouzée, A. *et al.* Imaging the electronic structure of valence orbitals in the XUV ionization of aligned molecules. *Journal of Physics B: Atomic, Molecular and Optical Physics* **47**, 124017 (2014).
14. Marceau, C. *et al.* Molecular Frame Reconstruction Using Time-Domain Photoionization Interferometry. *Phys. Rev. Lett.* **119**, 083401 (2017).
15. Thomann, I. *et al.* Direct Measurement of the Angular Dependence of the Single-Photon Ionization of Aligned N₂ and CO₂. *The Journal of Physical Chemistry A* **112**, 9382–9386 (2008).